

WASP-7: The brightest transiting-exoplanet system in the Southern hemisphere.

Coel Hellier¹, D.R. Anderson¹, M. Gillon², T.A. Lister^{1,3}, P.F.L. Maxted¹, D. Queloz², B. Smalley¹, A. Triaud², R.G. West⁴, D.M. Wilson¹, K. Alsubai⁵, S.J. Bentley¹, A. Collier Cameron⁵, L. Hebb⁵, K. Horne⁵, J. Irwin⁶, S.R. Kane⁷, M. Mayor², F. Pepe², D. Pollacco⁸, I. Skillen⁹, S. Udry², P.J. Wheatley¹⁰, D.J. Christian⁸, R. Enoch^{11,5}, C.A. Haswell¹¹, Y.C. Joshi⁸, A.J. Norton¹¹, N. Parley¹¹, R. Ryans⁸, R.A. Street^{3,8}, I. Todd⁸

ABSTRACT

We report that a Jupiter-mass planet, WASP-7b, transits the $V = 9.5$ star HD 197286 every 4.95 d. This is the brightest discovery from the WASP-South transit survey and the brightest transiting-exoplanet system in the Southern hemisphere. WASP-7b is among the densest of the known Jupiter-mass planets, suggesting that it has a massive core. The planet mass is $0.96^{+0.12}_{-0.18} M_{\text{Jup}}$, the radius $0.915^{+0.046}_{-0.040} R_{\text{Jup}}$ and the density $1.26^{+0.25}_{-0.21} \rho_{\text{Jup}}$.

Subject headings: planetary systems: individual: WASP-7b — stars: individual: HD 197286

1. Introduction

Transiting exoplanets are valuable discoveries since they offer the most opportunities for parametrization and study. The WASP project

¹Astrophysics Group, Keele University, Staffordshire, ST5 5BG, UK

²Observatoire de Genève, 51 ch. des Maillettes, 1290 Sauverny, Switzerland

³Las Cumbres Observatory, 6740 Cortona Dr. Suite 102, Santa Barbara, CA 93117, USA

⁴Department of Physics and Astronomy, University of Leicester, Leicester, LE1 7RH, UK

⁵School of Physics and Astronomy, University of St. Andrews, North Haugh, Fife, KY16 9SS, UK

⁶Department of Astronomy, Harvard University, 60 Garden Street, MS 10, Cambridge, Massachusetts 02138, USA

⁷Michelson Science Center, Caltech, MS 100-22, 770 South Wilson Avenue, Pasadena, CA 91125, USA

⁸Astrophysics Research Centre, School of Mathematics & Physics, Queen's University, University Road, Belfast, BT7 1NN, UK

⁹Isaac Newton Group of Telescopes, Apartado de Correos 321, E-38700 Santa Cruz de la Palma, Tenerife, Spain

¹⁰Department of Physics, University of Warwick, Coventry, CV4 7AL, UK

¹¹Department of Physics and Astronomy, The Open University, Milton Keynes, MK7 6AA, UK

(Pollacco et al. 2006) is one of a number of wide-area surveys, along with HAT (Bakos et al. 2002), TrES (O'Donovan et al. 2006) and XO (McCullough et al. 2005), all aimed at finding exoplanets transiting relatively bright stars, where they are easiest to observe. A prime aim is to fill out diagrams such as the exoplanet mass–radius plot, which has the potential to be a diagnostic tool for exoplanets comparable to the Hertzsprung–Russell diagram for stars.

WASP is the only one of the above surveys operating in both hemispheres. We report here on WASP-7 (= HD 197286), a new discovery from WASP-South that, at magnitude 9.5, is the brightest transiting-exoplanet system in the Southern hemisphere, being three magnitudes brighter than the brightest previously announced, WASP-4 (Wilson et al. 2008) and WASP-5 (Anderson et al. 2008). WASP-7b is also among the densest known Jupiter-mass exoplanets, extending the populated region of the mass–radius plot.

2. Observations

The WASP-South survey is described in Pollacco et al. (2006) and Wilson et al. (2008), while a discussion




Fig. 1.— The WASP-South lightcurve of WASP-7 folded on the 4.95-d transit period. The lower panel shows the transit phases enlarged, together with the best-fitting model.

Table 1: CORALIE radial velocities for WASP-7.

Time BJD – 245 0000	Rad Vel (km s ⁻¹)	σ_{RV} (km s ⁻¹)	Bis Span (km s ⁻¹) ^a
4329.71524	-29.8238	0.0214	-0.344
4330.79865	-29.7059	0.0182	-0.558
4359.54361	-29.8400	0.0196	-0.703
4362.52524	-29.9620	0.0202	-0.529
4364.55740	-29.7575	0.0221	-0.691
4376.57881	-29.8540	0.0171	-0.669
4378.59076	-29.8662	0.0194	-0.736
4379.57799	-29.7858	0.0222	-0.547
4380.63008	-29.8169	0.0173	-0.571
4382.66066	-29.9355	0.0210	-0.641
4386.61194	-29.8845	0.0229	-0.673

^a Bisector spans; $\sigma_{BS} \approx 2\sigma_{RV}$




Fig. 2.— The CORALIE radial-velocity curve for WASP-7 along with the best-fitting model (including the predicted Rossiter–McLaughlin effect, which is large for this fast rotator). The lower panel shows the bisector spans.

of our candidate selection methods can be found in Collier Cameron et al. (2007a), Pollacco et al. (2007), and references therein.

WASP-7 (= HD 197286) is a $V = 9.5$ F5V star in Microscopium. It was observed by WASP-South from May to mid-October in both 2006 and 2007, being recorded in two overlapping cameras, each an 11.1-cm-aperture Canon 200-mm f/1.8 lens backed by a 2k×2k e2V CCD. Exposure times were 30 s, with a typical cadence of 8 minutes. We obtained 5800 photometric data points from each camera in 2006, and a further 5700 data points from each in 2007.

The WASP-South lightcurves revealed dips with a depth of 0.007 magnitudes recurring with a 4.95-d period (Fig. 1). Spectroscopic observations were then obtained using the CORALIE spectrograph on the Euler 1.2-m telescope. Eleven radial-velocity measurements were obtained during 2007 Sept 15 to Oct 12 (Fig. 2; Table 1), establishing WASP-7b as a Jupiter-mass companion.

We used a line-bisector analysis to look for asymmetries in the spectral line profiles, as could be caused by contamination from an unresolved eclipsing binary (Queloz et al. 2001). Such a binary would produce bisector spans that vary in phase with the photometric period with an amplitude comparable to the radial-velocity amplitude. This is not seen in our data, supporting the conclusion that the radial-velocity variations are caused by a planet.

3. WASP-7 (HD 197286) parameters

The CORALIE spectra, when co-added, give a signal-to-noise ratio of ~ 50 in 0.01 Å bins, which

is suitable for a preliminary photospheric analysis of WASP-7.

We analysed the spectra using the UCLSYN package and ATLAS9 models, without convective overshooting (Castelli et al. 1997), leading to the parameters in Table 2. The effective temperature (T_{eff}) comes from an analysis of the H α line while the surface gravity ($\log g$) comes from the Na I D and Mg I b lines. An estimate of the microturbulence (ξ_t) comes from several clean and unblended Fe I and Fe II lines, while the ionization balance between Fe I and Fe II was used as an additional diagnostic of T_{eff} and $\log g$.

In addition to the spectral analysis, we have also used TYCHO, DENIS and 2MASS magnitudes to estimate the effective temperature using the Infrared Flux Method (Blackwell & Shallis 1977). This gives $T_{\text{eff}} = 6370 \pm 150$ K, which is in close agreement with that obtained from the spectroscopic analysis. These results are consistent with the F5V spectral type determined by Houk (1982).

The rotation rate of WASP-7 is $v \sin i = 17 \pm 2$ km s $^{-1}$. This is in line with expectations for an F5V star (Gray 1992 quotes $\langle v \sin i \rangle = 20$), but is the largest known among host stars of transiting planets, and leads to the prediction of a large Rossiter–McLaughlin effect of 98 ± 19 m s $^{-1}$ (e.g. Gaudi & Winn 2007).

We have investigated whether the star might be chromospherically active, however there is no sign of Ca H+K line emission in the CORALIE spectra. We have also looked for variability in the WASP data at the predicted rotation period of 3.7 ± 0.5 d, but find no such variability with an upper limit of 0.02 mag.

The Li I 6708 Å line is not detected in the CORALIE spectra ($\text{EW} \ll 1 \text{ mÅ}$, allowing us to derive an upper-limit on the lithium abundance of $\log n(\text{Li}/\text{H}) + 12 \ll 1.0$). However, the T_{eff} of this star implies that it is in the lithium gap (Böhm-Vitense 2004) and thus the lithium line does not provide an age constraint.

3.1. Planetary parameters

The CORALIE radial-velocity measurements were combined with the WASP-South photometry in a simultaneous Markov-chain Monte-Carlo (MCMC) analysis to find the parameters

Table 2: Parameters of WASP-7/HD 197286.

R.A. = 20 ^h 44 ^m 10.22 ^s , Dec. = −39°13′30.9″		
T_{eff}	6400 ± 100	K
$\log g$	4.3 ± 0.2	
[Fe/H]	0.0 ± 0.1	
ξ_t	1.5 ± 0.2	km s $^{-1}$
$v \sin i$	17 ± 2	km s $^{-1}$
Spectral Type	F5V	
Distance	140 ± 15	pc
V mag	9.51	

Table 3: System parameters for WASP-7.

Parameter	Value	
P (days)	4.954658	+0.000055 -0.000043
T_C (HJD)	2453985.0149	+0.0009 -0.0012
T_{14} (days) ^a	0.1573	+0.0024 -0.0018
R_P^2/R_*^2	0.00579	+0.00013 -0.00026
$b \equiv a \cos i/R_*$ ($R_\odot$)	0.08	+0.17 -0.08
i (deg)	89.6	+0.4 -0.9
e	0 (adopted)	
K_1 (km s^{-1})	0.097	+0.013 -0.013
γ (km s^{-1})	-29.8506	+0.0017 -0.0016
M_* ($M_\odot$)	1.28	+0.09 -0.19
R_* ($R_\odot$)	1.236	+0.059 -0.046
T_{eff} (K)	6400	± 100
$\log g_*$ (cgs)	4.363	+0.010 -0.047
M_P (M_J)	0.96	+0.12 -0.18
R_P (R_J)	0.915	+0.046 -0.040
ρ_P (ρ_J)	1.26	+0.25 -0.21
a (AU)	0.0618	+0.0014 -0.0033
$\log g_P$ (cgs)	3.421	+0.067 -0.071
T_P (K)	1379	+35 -23

^a T_{14} : duration, time from 1st to 4th contact.
Errors are 1- σ




Fig. 3.— The mass-radius diagram for Jupiter-mass transiting exoplanets. The WASP planets (points) are labelled by their number (data from the WASP consortium, papers in preparation). The remaining data (crosses) are taken from Pont (2008).

of the WASP-7 system. This process is described in detail in Collier Cameron et al. (2007b) and Pollacco et al. (2007). The optimal MCMC parameters are listed in Table 3.

An initial run found an eccentricity, $e < 0.17$, and we then adopted $e = 0$ for the solution presented here. In order to balance the weights of the photometry and radial velocities in the MCMC analysis we added a systematic error of 20 m s^{-1} to the radial velocities (as might arise, for example, from stellar activity) to reconcile χ^2 with the number of degrees of freedom.

WASP-7b is among the densest of the Jupiter-mass planets (Fig. 3), which suggests that it has a large core. The calculations presented by Fortney, Marley & Barnes (2007) suggest a core mass of ≈ 100 Earth masses, or ≈ 0.3 of the planet, with some dependence on the planetary age. Overall, WASP-7b adds to the finding that Hot Jupiters show a wide disparity in densities, from the denser WASP-5b (Anderson et al. 2008) and WASP-7b to the bloated TrES-4 (Mandushev et al. 2007) and WASP-12b (Hebb et al. 2008).

4. Acknowledgments

The WASP consortium comprises the Universities of Keele, Leicester, St. Andrews, the Queen's University Belfast, the Open University and the Isaac Newton Group. WASP-South is hosted by the South African Astronomical Observatory and we are grateful for their support and assistance. Funding for WASP comes from consortium universities and from the UK's Science and Technology Facilities Council.

REFERENCES

- Anderson D. R., et al., 2008, MNRAS, in press (arXiv:0801.1685)
- Bakos G. À., Lázár J., Papp I., Sári P., Green E. M., 2002, PASP, 114, 974
- Böhm-Vitense E., 2004, AJ, 128, 2435
- Blackwell D.E., Shallis M.J., 1977, MNRAS, 180, 177
- Castelli F., Gratton R. G., Kurucz R. L., 1997, A&A, 318, 841
- Collier Cameron A., et al., 2007a, MNRAS, 375, 951
- Collier Cameron A., et al., 2007b, MNRAS, 380, 1230
- Fortney J. J., Marely M. S., Barnes, J. W., 2007, ApJ, 659, 1661
- Gaudi B. S., Winn J. N., 2007, ApJ, 655, 550
- Gray D.F., 1992, The observation and analysis of stellar photospheres, 2nd edn, CUP, Cambridge
- Hebb L. et al., 2008, in preparation
- Houk N., 1982, Catalogue of two-dimensional spectral types for the HD stars, Vol. 3, Michigan Spectral Survey, Ann Arbor, Dep. Astron., Univ. Michigan.
- Mandushev, G. et al., 2007, ApJ, 667, L195
- McCullough P. R., Stys J., Valenti J., Fleming S., Janes K., Heasley J., 2005, PASP, 117, 783
- O'Donovan F. T., et al., 2006, ApJ, 651L, 610
- Pollacco D., et al., 2006, PASP, 118, 1407
- Pollacco D., et al., 2008, MNRAS, 385, 1576
- Pont F., 2008, <http://www.inscience.ch/transits/>
- Queloz D., et al., 2001, A&A, 379, 279
- Wilson D. M., et al., 2008, ApJ, 675, L113